

## Pneumoconiosis deaths

Health Priority D: Environmental and Occupational Health Hazards

Objective D2d: By 2010, reduce occupational pneumoconiosis illness and death by 30 percent below the 2000 baseline.

2010 Target: 0.3 deaths per 100,000 population

Indicator: Pneumoconiosis deaths

### Wisconsin Deaths With Pneumoconiosis as the Underlying or Contributing Cause of Death, by Sex and Age

Year	Total	Deaths per 100,000 Population	Number				
			Males	Females	Under Age 65	Ages 65-74	Age 75+
2000	21	0.4	21	0	3	3	15
2001	27	0.5	26	1	4	9	14
2002	26	0.5	25	1	4	8	14
2003	32	0.5	29	3	4	5	23
2004	33	0.6	32	1	5	7	21

Source: Resident death certificates, Bureau of Health Information and Policy, Division of Public Health, Department of Health and Family Services.

Note: Rates (deaths per 100,000 population) have been age-adjusted to the 2000 U.S. standard population. (See data documentation.)

### Wisconsin Deaths With Pneumoconiosis as the Underlying or Contributing Cause of Death, by Race/Ethnicity

Year	Total	African American*	American Indian*	Asian*	Hispanic	White*
2000	21	1	1	0	0	19
2001	27	3	0	0	0	24
2002	26	3	0	0	0	23
2003	32	2	0	0	0	30
2004	33	4	1	0	0	28

Source: Resident death certificates, Bureau of Health Information and Policy, Division of Public Health, Department of Health and Family Services.

\* Non-Hispanic